ROBINSON MEMORIAL HOSPITAL – RAVENNA, OHIO PHYSICIAN'S DIRECTIONS

	(IMPF	RINT PATIENT'S PLATE HERE)
DRUG ALLERGI	ES:	INITIAL HEIGHT Ib PREGNANT LACTATING
DATE	TIME	ORDER & SIGNATURE
		VTE PROPHYLAXIS & ANTICOAGULATION ORDER
		Labs prior to pharmacologic intervention (unless already ordered): BMP and CBC
		2. Pharmacologic DVT/VTE Prophylaxis: Dalteparin (Fragmin) 5000 units SQ every 24 hours Heparin 5000 units SQ every 8 hours Heparin 5000 units SQ every 12 hours Fondaparinux (Arixtra) 2.5 mg daily (only for ortho hip and knee)
		3. Compression Devices for DVT/VTE Prophylaxis: Sequential Compression Devices (SCDs) Plantar Pneumatic compression (Foot Pumps) Graduated compression stockings (Thigh-high TEDS)
		 4. Full anticoagulation/ Treatment dose: Acute Coronary Syndrome (UA/NSTEMI) Enoxaparin 1mg/kg SQ every 12 hours Enoxaparin 1mg/kg SQ every 24 hours (if patient's creatinine clearance is less than 30 mL/min) Heparin infusion: see Heparin weight-based protocol
		DVT/PE Dalteparin (Fragmin) 200 units/kg SQ every 24 hours (Maximum of 18,000 units per dose - if patient's weight is greater than 150kg, consider Heparin infusion) Heparin infusion: see Heparin weight-based protocol (consider if patient's creatinine clearance is less than 30 mL/min)
·		Atrial Arrhythmias Dalteparin (Fragmin) 100 units/kg SQ every 12 hours (Maximum of 10,000 units per dose - if patient's weight is greater than 150kg, consider Heparin infusion) Heparin infusion: see Heparin weight-based protocol (consider if patient's creatinine clearance is less than 30 mL/min)
·		Physician signature: Date/Time:

UNLESS CHECKED, ANOTHER BRAND OF DRUG IDENTICAL IN FORM AND CONTENT MAY BE DISPENSED.

FORM # 085241 (11/09)